



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

0 075 267
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: 82108507.3

22 Date of filing: 15.09.82

51 Int. Cl.³: **C 07 D 487/04, C 07 D 237/04,**
A 01 N 43/64
//
(C07D487/04, 249/00, 243/00),
(C07D487/04, 249/00, 237/00)

30 Priority: 19.09.81 JP 148443/81
19.02.82 JP 26512/82

43 Date of publication of application: 30.03.83
Bulletin 83/13

84 Designated Contracting States: CH DE FR GB IT LI NL


71 Applicant: **SUMITOMO CHEMICAL COMPANY, LIMITED,**
15 Kitahama 5-chome Higashi-ku, Osaka-shi Osaka-fu
(JP)

72 Inventor: Nagano, Eiki, 4-1-401, Ryodo-cho Nishinomiya,
Hyogo (JP)
Inventor: Hashimoto, Shunichi, 2-10-4-442,
Sonehigashi-machi, Toyonaka Osaka (JP)
Inventor: Yoshida, Ryo, 5-19, Azajizoyama Higashineno,
Kawanishi Hyogo (JP)
Inventor: Matsumoto, Hiroshi, 2-10-2-211,
Sonehigashi-machi, Toyonaka Osaka (JP)
Inventor: Kamoshita, Katsuzo, 2-3-11, Kofudai
Toyono-cho, Toyono-gun Osaka (JP)

74 Representative: **Vossius Vossius Tauchner Heunemann**
Rauh, Siebertstrasse 4 P.O. Box 86 07 67,
D-8000 München 86 (DE)

54 4-(2-Fluoro-4-halo-5-substituted phenyl)urazols, and their production and use.

57 A condensed phenylurazol of the formula:



wherein X is a chlorine atom or a bromine atom, Y and Z are each an oxygen atom or a sulfur atom and R is a C₁-C₃ alkyl group, an allyl group or a propargyl group and n is an integer of 4 or 5, which is useful as herbicide.

EP 0 075 267 A1

ACTORUM AG

BEST AVAILABLE COPY

0075267

SIEBERTSTRASSE 4 · 8000 MÜNCHEN 86 · PHONE: (089) 47 40 75
CABLE: BENZOLPATENT MÜNCHEN · TELEX 5-29 453 VOPAT D

SUMITOMO CHEMICAL COMPANY

Case: 603601

- 1 -

September 15, 1982


Our Ref.: S 098 EP

4-(2-FLUORO-4-HALO-5-SUBSTITUTED PHENYL)URAZOLS,

AND THEIR PRODUCTION AND USE

The present invention relates to 4-(2-fluoro-4-halo-5-substituted phenyl)urazols bearing a condensed hexahydropyridazine or hexahydrodiazepine ring (hereinafter referred to as "the condensed phenylurazol(s)"), and their production and use.

The said condensed phenylurazols are representable by the formula:



wherein X is a chlorine atom or a bromine atom, Y and Z are each an oxygen atom or a sulfur atom, R is a C_1 - C_3 alkyl group, an allyl group or a propargyl group and n is an integer of 4 or 5.

It is known that some 4-phenylurazols exhibit a herbicidal activity. For instance, U.S. patent 4,249,934, Japanese Patent Publication (unexamined) No. 44587/1978, etc. disclose that 2-(4-chlorophenyl-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]pyridazine-1,3(2H)-dione, 2-(4-chloro-2-fluorophenyl)-5,6,7,8-tetrahydro-1H-[1.2.4]-triazolo[1.2-a]pyridazine-1,3(2H)-dione, 2-(4-chlorophenyl)-1H,5H-[1.2.4]triazolo[1.2-a][1.2]diazepin-1,3(2H)-dione, etc. show a herbicidal activity. However, their

0075267

herbicidal effect is not always satisfactory.

It has been found that the condensed phenylurazols (I) show a strong herbicidal activity against a wide variety of weeds including Gramineaceous weeds, Cyperaceae weeds and broad-leaved weeds at small doses and do not produce any material phytotoxicity on various agricultural crops (e.g. corn, soybean, cotton, wheat). Examples of Gramineaceous weeds against which the condensed phenylurazols (I) show a herbicidal activity are barnyardgrass (Echinochloa crus-galli), green foxtail (Setaria viridis), large crabgrass (Digitaria sanguinalis), Johnsongrass (Sorghum halepense), wild oat (Avena fatua), black grass (Alopecurus myosuroides), goosegrass (Eleusine indica), annual bluegrass (Poa annua), bermudagrass (Cynodon dactylon), quackgrass (Agropyron repens), etc. Examples of broad-leaved weeds are tall morningglory (Ipomoea purpurea), velvetleaf (Abutilon theophrasti), sicklepod (Cassia obtusifolia), wild sunflower (Helianthus annus), cocklebur (Xanthium pensylvanicum), wild mustard (Brassica kaber), common chickweed (Stellaria media), common purslane (Portulaca oleracea), jimsonweed (Datura stramonium), hemp sesbania (Sesbania exaltata), sun spurge (Euphorbia helioscopia), black nightshade (Solanum nigrum), prickly sida (Sida spinosa), common ragweed (Ambrosia artemisiifolia), smartweed sp. (Polygonum sp.), redroot pigweed (Amaranthus retroflexus), bedstraw (Galium aparine), pineappleweed (Matricaria spp.), birdseye speedwell (Veronica persica), wild buckwheat (Polygonum


0075267

convolvulus), beggarticks (Bidens spp.), common lambs-quarters (Chenopodium album), bindweed (Calystegia japonica), monochoria (Monochoria vaginalis), Dopatrium junceum, waterwort (Elatine triandra), false pimpernel (Lindernia procumbens), toothcup (Rotala indica), arrowhead (Sagittaria pygmaea), etc. Examples of Cyperaceae weeds are nutsedge sp. (Cyperus microiria), purple nutsedge (Cyperus rotundus), yellow nutsedge (Cyperus esculentus), hardstem bulrush (Scirpus juncoides), nutsedge (Cyperus serotinus), water chestnut (Eleocharis kuroguwai), slender spikerush (Eleocharis acicularis), etc. Accordingly, the condensed phenylurazols (I) can be used as herbicides applicable to paddy field as well as agricultural plowed field. They are also useful as herbicides to be employed for crop field, orchard, tea garden, mulberry field, rubber plantation, forest, lawn, pasture, non-agricultural field, etc.

The condensed phenylurazols (I) can be produced by various procedures, among which typical examples are shown below:

Procedure A


The condensed phenylurazols of the formula (I) wherein at least one of Y and Z is an oxygen atom can be produced by reacting a phenylurazol of the formula:



(II)

0075267

wherein X, Y, Z and R are each as defined above but at least one of Y and Z is an oxygen atom with a dihaloalkane of the formula:




wherein A and A' are each a halogen atom (e.g. chlorine, bromine) and n is as defined above, usually in an inert solvent (e.g. dimethylformamide, dimethylsulfoxide, acetonitrile, ethylene glycol monomethyl ether, tetrahydrofuran, diethyl ether) at a temperature of -80 to 200°C.

Preferably, the phenylurazol (II) is previously metallized, and the metallized phenylurazol is subjected to reaction with the dihaloalkane (III). The metallization may be achieved by treatment of the phenylurazol (II) with a metallizing agent (e.g. sodium alkoxide, sodium hydride, n-butyl lithium) in an inert solvent (e.g. methanol, ethanol, diethyl ether, tetrahydrofuran) at a temperature of -80 to 160°C. The thus produced metallized phenylurazol can be subjected to reaction with the dihaloalkane (III) as above with or without its isolation from the reaction mixture.

Procedure B

The condensed phenylurazols of the formula (I) wherein Z is an oxygen atom and n is an integer of 4 can be produced by treatment of a urea of the formula:

0075267




(IV)

wherein R' is a lower alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl) and X, Y and R are each as defined above with a base (e.g. sodium methoxide, sodium hydroxide, potassium hydroxide) in an inert solvent (e.g. water, .. toluene, benzene, methanol, ethanol) at a temperature of 0 to 200°C.

Procedure C

The condensed phenylurazols of the formula (I) wherein Z is an oxygen atom and n is an integer of 4 can be produced by reacting a urea of the formula:




(V)

wherein X, R and Y are each as defined above with phosgene in an inert solvent (e.g. toluene, tetrahydrofuran, 1,4-dioxane) at a temperature of from 0°C to the boiling temperature of the solvent. When desired, a base (e.g. triethylamine, pyridine, diethylaniline) may be present in the reaction system.

Procedure D

The condensed phenylurazols of the formula (I) wherein Z is a sulfur atom and n is an integer of 4 can be produced by reacting a urea of the formula:

0075267



(V)


wherein X, R and Y are each as defined above with carbon disulfide in an inert solvent (e.g. methanol, ethanol, tetrahydrofuran) in the presence of a base such as an alkali metal hydroxide (e.g. sodium hydroxide, potassium hydroxide) or a tertiary amine (e.g. pyridine, triethylamine).

Procedure E

The condensed phenylurazols of the formula (I) wherein Y and Z are each a sulfur atom can be produced by reacting the corresponding condensed phenylurazol of the formula (I) wherein Y and Z are each an oxygen atom with phosphorus pentasulfide in an inert solvent (e.g. xylene, toluene, hexane) at a boiling temperature of the solvent.

The production of the starting materials is summarized in the following scheme:

0075267




wherein X, Y, Z, R and R' are each as defined above.

Namely, the phenol (VI) is selectively nitrated to the nitrophenol (VII), which is subjected to alkylation, alkenylation or alkynylation. The resultant nitrobenzene (VIII) is subjected to reduction, and the resulting aniline (IX) is reacted with phosgene or thiophosgene to give the isocyanate or isothiocyanate (X), which is reacted with an N-alkoxycarbonylhexahydropyridazine to give the urea (IV). Hydrolysis and decarboxylation of the urea (IV) give the urea (V). The above aniline (IX) is reacted with a 2-alkoxycarbonylhydrazinecarbonyl chloride, followed by

0075267

cyclization with a base to give the phenylurazol (II). Alternatively, the phenylurazol (II) can be produced by reacting the isocyanate or isothiocyanate (X) with an alkoxycarbonylhydrazine, followed by cyclization with a base.

Explaining the above conversions more in detail, the nitration of the phenol (VI) to the nitrophenol (VII) may be accomplished by application of a per se conventional nitration procedure to the former. Usually, however, the indirect nitration which consists of the following three steps is favorable in achievement of the selective nitration at the desired position:



wherein X is as defined above. Thus, the phenol (VI) (Finger et al.: J.Am.Chem.Soc., 81, 94 (1959)) is converted into its alkali metal salt by treatment with an aqueous solution of an alkali metal hydroxide (e.g. sodium hydroxide, potassium hydroxide), and the resulting salt is reacted with an alkyl haloformate such as methyl chloro-

0075267

formate in water at a temperature of 0 to 10°C. The thus prepared carbonic ester (XI) is nitrated with a mixture of conc. sulfuric acid and conc. nitric acid at room temperature. Then, the nitrobenzene (XII) thus obtained is hydrolyzed with an aqueous alkaline solution such as an aqueous sodium hydroxide solution at a temperature of 40 to 80°C to give the nitrophenol (VII).

The alkylation, alkenylation or alkynylation for conversion of the nitrophenol (VII) into the nitrobenzene (VIII) may be carried out by treatment of the former with an alkali metal carbonate (e.g. potassium carbonate), an alkali metal hydride (e.g. sodium hydride) or an alkali metal alkoxide (e.g. sodium methoxide) and reacting the resultant alkali metal salt with a halide of the formula: R-Q wherein Q is a halogen atom (e.g. chlorine, bromine, iodine) and R is as defined above in a polar solvent (e.g. water, dimethylformamide, acetonitrile, acetone, dimethylsulfoxide), usually at a temperature of 10 to 200°C, preferably of 30 to 100°C. The use of a phase transfer catalyst such as tetrabutylammonium bromide is favorable for smooth accomplishment of the reaction.

Reduction of the nitrobenzene (VIII) to the aniline (IX) may be achieved in various procedures. When, for instance, R in the compound (VIII) is C₁-C₄ alkyl, there may be adopted a per se conventional reduction procedure for converting a nitro group into an amino group wherein a reducing agent such as sodium sulfide or iron powder or

0075267

catalytic reduction is employed. One of typical procedures comprises introduction of a 3 molar amount of hydrogen into a reaction system comprising one molar amount of the compound (VIII) and a 1/10 to 1/100 molar amount of platinum dioxide at room temperature under atmospheric pressure. Another typical procedure comprises admixing an acetic acid solution containing one molar amount of the compound (VIII) with a 5 % acetic acid solution containing a 2.5 to 5.0 molar amount of iron powder such as reductive iron or electrolytic iron and effecting the reaction at a temperature of 80 to 100°C. When R in the compound (VIII) is propargyl or allyl, there may be adopted reduction with iron powder. For instance, an acetic acid solution containing one molar amount of the compound (VIII) may be admixed with a 5 % acetic acid solution containing a 2.5 to 5.0 molar amount of iron powder such as reductive iron or electrolytic iron at a temperature of 80 to 120°C, preferably of 90 to 110°C, for a period of 0.5 to 5 hours.

The aniline (IX) is converted into the isocyanate or isothiocyanate (X) by reacting the former with phosgene or thiophosgene in an inert solvent (e.g. toluene, benzene, ethyl acetate, tetrahydrofuran, 1,4-dioxane).


The isocyanate or isothiocyanate (X) can be converted into the urea (IV) by reacting the former with an N-alkoxycarbonylhexahydropyridazine in an inert solvent (e.g. benzene, toluene, hexane, tetrahydrofuran), preferably in the presence of an organic base (e.g. pyridine, triethyl-

0075267

amine), at a temperature from room temperature (ca. 20°C) to the boiling temperature of the solvent.

Hydrolysis and decarboxylation of the urea (IV) by treatment with an aqueous alkaline solution at a temperature of 0 to 100°C affords the urea (V).

Alternatively, the phenylurazol (II) can be produced by reacting the isocyanate or isothiocyanate (X) with an alkoxy-carbonylhydrazine or alkoxythiocarbonylhydrazine in an inert solvent (e.g. toluene, chloroform, tetrahydrofuran) at a temperature of 0 to 100°C, followed by treatment of the resultant semicarbazide of the formula:



wherein R'' is an alkyl group and X, Y, Z and R are each as defined above with an aqueous alkaline solution such as potassium hydroxide solution and then with a mineral acid (e.g. hydrochloric acid, sulfuric acid).

The above obtained aniline (IX) can be converted into the semicarbazide (XIII) by reacting the former with a 2-alkoxycarbonylhydrazinecarbonyl chloride in an inert solvent (e.g. benzene, toluene, diethyl ether, tetrahydrofuran, dioxane) at the boiling temperature of the solvent.

The above produced condensed phenylurazols (I) and their intermediates are, when desired, purified by a per se conventional procedure such as recrystallization or column chromatography.

0075267

Practical and presently preferred embodiments for production of the condensed phenylurazols (I) are shown in the following Examples.

Example 1

To a solution of metallic sodium (0.7 g) in methanol (40 ml), 4-(4-chloro-2-fluoro-5-methoxy)urazol (3.87 g) was added, and the resultant mixture was concentrated under reduced pressure. A solution of 1,4-dibromobutane (3.5 g) in dimethylformamide (40 ml) was added thereto. The resulting mixture was stirred at 100 to 110°C for 4 hours. After allowed to cool to room temperature, water was added to the mixture, which was then extracted with ether. The ether layer was washed with water, dried and concentrated to give crystals. The crystals were washed with ether to give 1.26 g of 2-(4-chloro-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]pyridazine-1,3(2H)-dione (Compound No. 1) as colorless crystals. M.P., 131 - 132°C. IR ν_{\max} (cm⁻¹): 1770, 1720. NMR δ (ppm): 1.9 (4H, m), 3.6 (4H, m), 3.85 (3H, s), 6.85 (1H, d, J=6Hz), 7.3 (1H, d, J=10Hz).

Example 2

To a solution of metallic sodium (0.46 g) in methanol (20 ml), 2-(4-chloro-2-fluoro-5-methoxyphenyl)-urazol (2.6 g) was added, and the resultant mixture was concentrated under reduced pressure. The residue was dissolved in dimethylformamide (20 ml), 1,5-dibromopentane (2.3 g) was added thereto, and the resulting mixture was

0075267

stirred at 100 to 120°C for 4 hours. After allowed to cool, water was added to the mixture, which was then extracted with ethyl acetate. The extract was washed with a saturated sodium chloride solution, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography and recrystallized from methanol to give 0.42 g of 2-(4-chloro-2-fluoro-5-methoxyphenyl)-1H,5H-[1.2.4]triazolo[1.2-a]-[1.2]diazepine-1,3(2H)-dione (Compound No. 3). M.P., 121.5 - 123°C. IR ν_{\max} (cm^{-1}): 1750, 1680. NMR δ (ppm): 1.8 (6H, m), 3.9 (3H, s).

Example 3

Ethyl 2-[(4-chloro-2-fluoro-5-(1-methylethoxy)-phenyl)aminothioxomethyl]-3,4,5,6-tetrahydro-1(2H)-pyridazinecarboxylate (0.66 g) was dissolved in toluene (20 ml), and several drops of sodium methoxide were added thereto, followed by heating under reflux for 3 hours. Water was added to the resultant mixture, which was then extracted with toluene. The extract was washed with a saturated sodium chloride solution, dried and concentrated to give 0.3 g of 2-[4-chloro-2-fluoro-5-(1-methylethoxy)-phenyl]-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]-pyridazine-3-thioxo-1(2H)-one (Compound No. 15). n_D^{25} 1.5720. IR ν_{\max} (cm^{-1}): 1740, 1490, 1250. NMR δ (ppm): 1.4 (6H, d), 2.0 (4H, m), 3.65 (2H, m), 4.0 (2H, m), 4.45 (1H, m), 6.9 (1H, d), 7.25 (1H, d).

Example 4


0075267

2-(4-Bromo-5-ethoxy-2-fluorophenyl)-1H,5H-[1.2.4]-
triazolo[1.2-a][1.2]diazepine-1,3(2H)-dione (0.63 g) was
dissolved in xylene (10 ml), and phosphorus pentasulfide
(0.46 g) was added thereto, followed by heating under reflux
for 8 hours. The reaction mixture was filtered to eliminate
insoluble materials, and the filtrate was concentrated under
reduced pressure. The residue was crystallized and washed
with methanol to give 0.65 g of 2-(4-bromo-5-ethoxy-2-
fluorophenyl)-1H,5H-[1.2.4]triazolo[1.2-a][1.2]diazepine-
1,3(2H)-dithione (Compound No. 9). M.P., 173.5 - 174.5°C.
IR ν_{\max} (cm^{-1}): 1290. NMR δ (ppm): 1.45 (3H, t), 1.9 (6H,
m), 4.07 (2H, q).

Some additional examples of the condensed
phenylurazol (I) produced in the same manner as above are
shown in Table 1.

0075267

Table 1



(I)

Compound No.	X	Y	Z	n	R	Physical property
1	Cl	O	O	4	-CH ₃	M.P. 131-132°C
2	Cl	S	O	4	-CH ₃	M.P. 146-147°C
3	Cl	O	O	5	-CH ₃	M.P. 121.5-123°C
4	Cl	S	S	5	-CH ₃	M.P. 229-231°C
5	Br	O	O	4	-CH ₃	M.P. 162.5-164.5°C
6	Cl	O	O	4	-CH ₂ CH ₃	M.P. 116-117°C
7	Br	O	O	4	-CH ₂ CH ₃	M.P. 141-143°C
8	Br	O	O	5	-CH ₂ CH ₃	M.P. 56-58°C
9	Br	S	S	5	-CH ₂ CH ₃	M.P. 173.5-174.5°C
10	Cl	O	O	4	-CH ₂ CH ₂ CH ₃	M.P. 97.5-99°C
11	Cl	S	S	4	-CH ₂ CH ₂ CH ₃	M.P. 181.5-182.5°C
12	Cl	O	O	5	-CH ₂ CH ₂ CH ₃	n _D ^{22.0} 1.5310
13	Br	O	O	4	-CH ₂ CH ₂ CH ₃	M.P. 92-93°C
14	Cl	O	O	4	-CH(CH ₃) ₂	M.P. 140-142.5°C
15	Cl	S	O	4	-CH(CH ₃) ₂	n _D ²⁵ 1.5720
16	Cl	S	O	4	-CH ₂ CH=CH ₂	M.P. 58-63°C
17	Br	O	O	4	-CH ₂ CH=CH ₂	M.P. 109-111°C
18	Cl	S	O	4	-CH ₂ C≡CH	M.P. 148-150°C

0075267


Example 5

A solution of 2-fluoro-4-chloro-5-n-propoxyphenyl isocyanate (1.8 g) in benzene (10 ml) was added to a toluene solution containing methoxycarbonylhydrazine (0.71 g), and the resultant mixture was allowed to stand overnight. The precipitated crystals were collected by filtration and dissolved in a 4M aqueous sodium hydroxide solution (10 ml) under heating. To the resulting solution, conc. hydrochloric acid was added, and the precipitated crystals were collected by filtration to give 2.6 g of 4-(4-chloro-2-fluoro-5-n-propoxyphenyl)urazol. M.P., 181 - 182°C. IR ν_{\max} (cm^{-1}): 3150, 1710, 1680. NMR δ (ppm): 1.1 (3H, t), 4.0 (3H, q), 7.0 (1H, d), 7.3 (1H, d)

Some examples of the phenylurazol (II) produced in the same manner as above are shown in Table 2.

0075267

Table 2



(II)

X	Y	Z	R	Physical property
Cl	O	O	-CH ₃	M.P. 223-224°C
Cl	S	O	-CH ₃	Glassy; IR ν_{\max} (cm ⁻¹): 3160, 1720
Cl	O	O	-CH ₂ CH ₂ CH ₃	M.P. 181-182°C
Br	O	O	-CH ₃	M.P. 229-230°C
Br	O	O	-CH ₂ CH ₃	M.P. 215-216°C
Br	O	O	-CH ₂ CH ₂ CH ₃	M.P. 189-189.5°C
Cl	O	O	-CH ₂ C=CH	M.P. 217-217.5°C

Example 6


4-Chloro-2-fluoro-5-(1-methylethoxy)phenyl isothiocyanate (2.8 g) was added to a solution of 2-ethoxycarbonyl-3,4,5,6-tetrahydro-(1H,2H)-pyridazine (1.8 g) and several drops of triethylamine in toluene (5 ml), and the resultant mixture was stirred at 25°C overnight. Water was added to the mixture, which was then extracted with toluene. The toluene layer was washed with water, dried and concentrated. The residue was purified by silica gel column chromatography to give 2 g of ethyl 2-(4-chloro-2-fluoro-5-(1-methylethoxy)phenylaminothioxomethyl)-3,4,5,6-tetrahydro-1(2H)pyridazine carboxylate. $n_D^{27.5}$ 1.5534. NMR δ (ppm): 1.3 (3H, t), 1.4 (6H, d), 1.8 (4H, m), 4.35 (2H, q),

0075267

7.1 (1H, d), 8.0 (1H, d), 8.45 (1H, m, $-\text{NH}$).

Some examples of the urea (IV) produced in the same manner as above are shown in Table 3.

Table 3



(IV)

X	Y	R	R'	Physical property
Cl	S	$-\text{CH}_3$	$-\text{CH}_2\text{CH}_3$	$n_D^{24.0}$ 1.5751
Cl	S	$-\text{CH}(\text{CH}_3)_2$	$-\text{CH}_2\text{CH}_3$	$n_D^{27.5}$ 1.5534
Cl	O	$-\text{CH}_2\text{C}\equiv\text{CH}$	$-\text{CH}_2\text{CH}_3$	M.P. 135.5-136.5°C
Cl	S	$-\text{CH}_2\text{C}\equiv\text{CH}$	$-\text{CH}_2\text{CH}_3$	M.P. 105-109°C
Br	S	$-\text{CH}_2\text{CH}_2\text{CH}_3$	$-\text{CH}_2\text{CH}_3$	Glassy
Br	S	$-\text{CH}_2\text{CH}=\text{CH}_2$	$-\text{CH}_2\text{CH}_3$	M.P. 80-86°C
Br	O	$-\text{CH}_2\text{CH}=\text{CH}_2$	$-\text{CH}_2\text{CH}_3$	M.P. 77-80°C
Cl	O	$-\text{CH}_2\text{CH}_2\text{CH}_3$	$-\text{CH}_2\text{CH}_3$	n_D^{25} 1.5313

Example 7

A solution of ethyl 2-(4-chloro-2-fluoro-5-(1-methylethoxy)phenylaminothioxomethyl)-3,4,5,6-tetrahydro-1(2H)pyridazinecarboxylate (0.8 g) in a 5 % ethanolic potassium hydroxide solution (10 ml) was heated under reflux for 3 hours. After allowed to cool to room temperature, ethanol was removed under reduced pressure. The residue was dissolved in ether, washed with water, dried over anhydrous magnesium sulfate and concentrated to give 0.3 g of 2-(4-chloro-2-fluoro-5-(1-methylethoxy)phenylaminothioxomethyl)-

0075267

3,4,5,6-tetrahydro-(1H,2H)-pyridazine. M.P., 113 - 114.5°C.
NMR δ (ppm): 1.35 (6H, d), 1.7 (4H, m), 2.95 (2H, m), 3.45 (1H, t), 4.2 (2H, m), 4.4 (1H, m), 6.95 (1H, d), 8.25 (1H, d), 9.85 (1H, m).

Some examples of the urea (V) produced in the same manner as above are shown in Table 4.

Table 4



(V)

X	Y	R	Physical property
Cl	S	-CH(CH ₃) ₂	M.P. 113-114.5°C
Cl	O	-CH ₂ CH ₂ CH ₃	M.P. 120-122.5°C
Cl	O	-CH ₂ C \equiv CH	M.P. 89.5-91.5°C
Cl	S	-CH ₂ C \equiv CH	M.P. 96.5-99°C
Cl	O	-CH ₂ CH=CH ₂	M.P. 91-95°C
Br	O	-CH ₂ CH=CH ₂	M.P. 80-86°C

Example 8


A solution of 4-chloro-2-fluoro-5-isopropoxy-aniline (30 g) in toluene (100 ml) was added to a 1 M phosgene/toluene solution (500 ml) at room temperature (ca. 20°C), followed by heating under reflux. The mixture was concentrated under reduced pressure, and the residue was distilled to give 26 g of 4-chloro-2-fluoro-5-isopropoxy-phenyl isocyanate as pale yellow crystals. M.P., 36 - 37°C. B.P., 90 - 91°C/3 mmHg.

0075267

IR ν_{\max} (cm^{-1}): 2240.

Some examples of the isocyanate or isothiocyanate (X) produced in the same manner as above are shown in Table 5.

Table 5



(X)

X	Y	R	Physical property
Cl	O	CH ₃ -	M.P. 42 - 44.5°C
Cl	O	n-C ₃ H ₇ -	M.P. 43 - 44°C
Cl	O	iso-C ₃ H ₇ -	M.P. 36 - 37°C
Br	O	C ₂ H ₅ -	M.P. 35 - 36.5°C
Cl	O	CH ₂ =CHCH ₂ -	B.P. 107°C/3 mmHg; n_D^{16} 1.5481
Cl	O	CH≡CCH ₂ -	M.P. 61.5 - 62.5°C
Cl	S	CH ₃ -	M.P. 79-80°C
Cl	S	CH≡CCH ₂ -	M.P. 105-109°C
Cl	S	CH ₂ =CHCH ₂ -	n_D^{24} 1.6228
Br	S	CH≡CCH ₂ -	M.P. 71-74°C
Br	S	CH ₂ =CHCH ₂ -	n_D^{24} 1.6421

Example 9

A mixture of 4-chloro-2-fluoro-5-isopropoxy-nitrobenzene (13.5 g) and platinum dioxide (0.4 g) in ethanol (300 ml) was subjected to catalytic reduction with hydrogen at room temperature (ca. 20°C) under atmospheric pressure, whereby a designed amount of hydrogen was

0075267

absorbed. The resultant mixture was filtered to remove insoluble materials, and the filtrate was concentrated. The residue was subjected to purification by silica gel chromatography to give 5.6 g of 4-chloro-2-fluoro-5-isopropoxyphenylaniline. $n_D^{24.5}$ 1.5360.

NMR ($CDCl_3$) δ (ppm): 1.3 (6H, d, $J=6\text{Hz}$), 3.7 (2H, m, $J=1.5\text{Hz}$), 4.35 (1H, q, $J=6\text{Hz}$), 6.45 (1H, d, $J=7\text{Hz}$), 7.1 (1H, d, $J=10\text{Hz}$).

IR ν_{max} (cm^{-1}): 3450, 3550.

Example 10

A suspension of electrolytic iron powder (3.5 g) in a 5 % aqueous acetic acid solution (5 ml) was heated to 90°C , and a solution of 4-chloro-2-fluoro-5-(2-propynyloxy)-nitrobenzene (5.7 g) in acetic acid (40 ml) was dropwise added thereto at the same temperature. The resultant mixture was stirred at $90 - 105^\circ\text{C}$ for 1 hour and allowed to cool to room temperature. Water (200 ml) was added thereto. Insoluble materials were filtered off, and the filtrate was neutralized, followed by extraction with ethyl acetate. The extract was dried over anhydrous magnesium sulfate and concentrated. The residue was washed with petroleum ether and carbon tetrachloride to give 3.6 g of 4-chloro-2-fluoro-5-(2-propynyloxy)aniline. M.P. $61.0 - 61.5^\circ\text{C}$.


NMR ($CDCl_3$) δ (ppm): 2.5 (1H, t, $J=2\text{Hz}$), 3.4 - 4.2 (2H, m, $J=16\text{Hz}$), 4.15 (2H, d, $J=2\text{Hz}$), 6.5 (1H, d, $J=8\text{Hz}$), 6.95 (1H, d, $J=10\text{Hz}$).

IR ν_{max} (cm^{-1}): 3460, 3360, 3280, 2100.

0075267

Some examples of the aniline (IX) produced in the same manner as above are shown in Table 6.

Table 6



(IX)

X	R	Physical property
Cl	C ₂ H ₅ -	n _D ^{24.5} 1.5503
Br	C ₂ H ₅ -	n _D ^{25.0} 1.5680
Cl	n-C ₃ H ₇ -	n _D ^{24.5} 1.5386
Br	n-C ₃ H ₇ -	n _D ^{26.0} 1.5618
Cl	iso-C ₃ H ₇ -	n _D ^{24.5} 1.5360
Br	iso-C ₃ H ₇ -	n _D ^{25.0} 1.5547
Cl	CH ₂ =CHCH ₂ -	n _D ¹⁹ 1.5598
Cl	CH≡CCH ₂ -	M.P. 61.0 - 61.5°C
Cl	CH≡C-CH- CH ₃	M.P. 67.0 - 68°C

Example 11

To a solution of 2-chloro-4-fluoro-5-nitrophenol (19.1 g) in acetonitrile (100 ml), there was added anhydrous potassium carbonate (8 g). After stirring at room temperature (ca. 20°C) for several minutes, isopropyl iodide (25 g) was added thereto, and the resultant mixture was heated under reflux for 3 hours. After allowed to cool to room temperature (ca. 20°C), water was added thereto, and the reaction mixture was extracted with ether. The ether

0075267


extract was washed with a 5 % aqueous sodium hydroxide solution and water in order, dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure, and the residue was recrystallized from ethanol to give 13.5 g of 4-chloro-2-fluoro-5-isopropoxy-nitrobenzene. M.P., 61.3 - 62.4°C.

NMR (CDCl_3) δ (ppm): 1.42 (6H, d, $J=7\text{Hz}$), 4.3 - 4.8 (1H, m), 7.28 (1H, d, $J=10\text{Hz}$), 7.48 (1H, d, $J=6\text{Hz}$).

Some examples of the nitrobenzene (VIII) produced in the same manner as above are shown in Table 7.

0075267

Table 7



(VIII)

X	R	Physical property
Cl	-CH ₃	M.P. 67.5 - 69.8°C
Br	-CH ₃	M.P. 72.2°C
Cl	-CH ₂ CH ₃	M.P. 47 - 48°C
Br	-CH ₂ CH ₃	M.P. 46 - 46.5°C
Cl	-CH ₂ CH ₂ CH ₃	M.P. 46 - 47°C
Br	-CH ₂ CH ₂ CH ₃	M.P. 46.8 - 47.4°C
Cl	-CH(CH ₃) ₂	M.P. 61.3 - 62.4°C
Br	-CH(CH ₃) ₂	M.P. 65.5 - 66.5°C
Cl	-CH(CH ₃)CH ₂ CH ₃	M.P. 59.6 - 60.6°C
Cl	-CH ₂ CH=CH ₂	n _D ^{17.0} 1.5601
Cl	-CHCH=CH ₂ CH ₃	M.P. 41.0 - 41.5°C
Cl	-CH ₂ C≡CH	M.P. 88 - 89°C
Cl	-CHC≡CH CH ₃	M.P. 87 - 88°C

Example 12

2-Chloro-4-fluorophenol (83.4 g) was added to a solution of sodium hydroxide (27.7 g) in water (450 ml), and methyl chloroformate (69.2 g) was dropwise added thereto at a temperature below 10°C. Precipitated crystals were collected by filtration and washed with water to give methyl

0075267

(2-chloro-4-fluorophenyl)formate (134.8 g). M.P., 69 - 71°C.

Methyl (2-chloro-4-fluorophenyl)formate (134.8 g) obtained above was suspended in conc. sulfuric acid (50 ml). To the suspension, a mixture of conc. sulfuric acid (50 ml) and conc. nitric acid (50 ml) was added at about 30°C, and the mixture was stirred at this temperature for 1 hour. The reaction mixture was poured into ice water, and precipitated crystals were collected and washed with water to give methyl (2-chloro-4-fluoro-5-nitrophenyl)formate (143 g). M.P., 50 - 55°C.

The product obtained as above was combined with sodium hydroxide (27 g) and water (300 ml), and the resultant mixture was refluxed for 4 hours. Precipitated insoluble materials were filtered using a celite, and the filtrate was acidified with conc. hydrochloric acid. Precipitated crystals were collected by filtration and washed with water to obtain 76.3 g of 2-chloro-4-fluoro-5-nitrophenol. M.P., 106 - 107°C.

NMR (CDCl_3 , D_6 -DMSO) δ (ppm): 7.25 (1H, d, $J=10\text{Hz}$), 7.64 (1H, d, $J=6\text{Hz}$).

IR $\nu_{\text{max}}^{\text{nujol}}$ (cm^{-1}): 3370.

Example 13

To a solution of 2-bromo-4-fluorophenol (28 g) in a solution of sodium hydroxide (7 g) in water (100 ml), methyl chloroformate was dropwise added thereto at a temperature below 10°C. The produced crystals were

0075267

collected by filtration and washed with water to give methyl 2-bromo-4-fluorophenylformate (41 g). M.P., 80.7°C.

The above product was suspended in conc. sulfuric acid (13 ml), a mixture of conc. sulfuric acid (13 ml) and conc. nitric acid (13 ml) was dropwise added thereto at about 30°C, and the resultant mixture of stirred for 30 minutes. The reaction mixture was poured into ice water. The produced crystals were collected by filtration and washed with water to give methyl 2-bromo-4-fluoro-5-nitroformate (38.3 g) as yellow crystals. M.P., 63.5 - 64.5°C.

The thus obtained product was admixed with sodium hydroxide (6.2 g) and water (100 ml) and heated under reflux for 3 hours. The insoluble materials were eliminated by filtration, and the filtrate was acidified with hydrochloric acid. The precipitated crystals were collected by filtration and washed with water to give 25 g of 2-bromo-4-fluoro-5-nitrophenol. M.P., 126 - 127°C.

NMR (CDCl_3 , D_6 -DMSO) δ (ppm): 7.42 (1H, d, $J=10\text{Hz}$), 7.65 (1H, d, $J=6\text{Hz}$).

IR $\nu_{\text{max}}^{\text{nujol}}$ (cm^{-1}): 3450.

In the practical usage of the condensed phenylurazols (I), they may be applied as such or in any composition form such as emulsifiable concentrates, wettable powders, suspensions, granules or fine granules.

The concentration of the active ingredient in such composition form is usually within a range of 0.1 to 90 % by weight, preferably of 1 to 80 % by weight.

0075267

For formulation of those compositions, a solid or liquid carrier or diluent may be used. As the solid carrier or diluent, there may be employed mineral powders (e.g. kaolin clay, bentonite, talc, diatomaceous earth, sericite, synthetic hydrated silicon dioxide). As the liquid carrier or diluent, there may be employed aromatic hydrocarbons (e.g. xylene, methylnaphthalene), ketones (e.g. cyclohexanone, isophorone), chlorobenzene, dimethylformamide, cello-solve, ethylene glycol, water, etc.

A surface active agent used for emulsification, dispersion or spreading may be any of the non-ionic, anionic, cationic and amphoteric type of agents. Examples of the surface active agent include polyoxyethylene alkyl ethers, polyoxyethylene alkylaryl ethers, polyoxyethylene fatty acid esters, sorbitan fatty acid esters, polyoxyethylene sorbitan fatty acid esters, oxyethyleneoxypropylene polymers, polyoxyethylene alkyl phosphates, fatty acid salts, alkyl sulfates, alkyl sulfonates, alkylaryl sulfonates, alkyl phosphates, polyoxyethylene alkyl sulfates, quaternary ammonium salts and the like. If necessary, gelatin, casein, sodium alginate, starch, agar, polyvinyl alcohol, ligninsulfonates, isopropyl acid phosphate, alginates or the like may be used as an auxiliary agent.

Practical embodiments of the herbicidal composition according to the invention are illustratively shown in the following examples wherein part(s) and % are by weight. The compound number of the active ingredient corresponds to

0075267

the one in Table 1.

Formulation Example 1

Eighty parts of Compound No. 10, 3 parts of alkyl-sulfate, 2 parts of ligninsulfonate and 15 parts of synthetic hydrated silicon dioxide are well mixed while being powdered to obtain a wettable powder.

Formulation Example 2

Twenty parts of Compound No. 14, 10 parts of polyoxyethylene alkylaryl ether, 50 parts of cyclohexanone and 20 parts of xylene are well mixed while being powdered to obtain an emulsifiable concentrate.

Formulation Example 3

0.1 Part of Compound No. 10, 1 part of synthetic hydrated silicon dioxide, 35 parts of bentonite and 63.9 parts of kaolin clay are well mixed while being powdered. The mixture is then kneaded with water, granulated and dried to obtain granules.

Formulation Example 4

Three parts of Compound No. 1, 0.3 part of isopropyl acid phosphate, 66.7 parts of kaolin clay and 30 parts of talc are well mixed while being powdered to obtain a dust.

Formulation Example 5

Eighty parts of Compound No. 3, 5 parts of polyoxyethylene alkylaryl ether and 15 parts of synthetic hydrated silicon dioxide are well mixed while being powdered to obtain a wettable powder.

0075267

Formulation Example 6

Ten parts of Compound No. 10, 7 parts of polyoxyethylene alkylaryl ether, 3 parts of alkylarylsulfate and 80 parts of cyclohexanone are well mixed while being powdered to obtain an emulsifiable concentrate.

Formulation Example 7

One part of Compound No. 9, 1 part of synthetic hydrated silicon dioxide, 5 parts of ligninsulfonate and 93 parts of kaolin clay are well mixed while being powdered. The mixture is then kneaded with water, granulated and dried to obtain granules.

Formulation Example 8

Three parts of Compound No. 3, 0.5 part of isopropyl acid phosphate, 66.5 parts of kaolin clay and 30 parts of talc are well mixed while being powdered to obtain a dust.

Formulation Example 9

Twenty parts of Compound No. 9 is mixed with 60 parts of an aqueous solution containing 3 % polyoxyethylene sorbitan monooleate and pulverized until the particle size of the active ingredient becomes less than 3 microns. Twenty parts of an aqueous solution containing 3 % of sodium alginate as a dispersing agent are incorporated therein to obtain a suspension.

The dosage rate of the condensed phenylurazols (I) may vary depending upon the application mode such as pre-emergence treatment or post-emergence treatment, etc.

007526

Generally, however, the dosage rate is from 0.1 to 50 grams, preferably from 0.2 to 30 grams, of the active ingredient per are. Besides, the condensed phenylurazols (I) of the invention may be used together with other herbicides to improve their activity as herbicides, and in some cases, a synergistic effect can be expected.

The application of the condensed phenylurazols (I) as herbicides will be illustratively shown in the following Examples wherein the phytotoxicity to crop plants and the herbicidal activity on weeds were evaluated as follows: the aerial parts of the test plants were cut off and weighed (fresh weight); the percentage of the fresh weight of the treated plant to that of the untreated plant was calculated with the latter fresh weight taken as 100; and the phytotoxicity and the herbicidal activity were evaluated by the standard given in the table below.

Rating value	Fresh weight (percentage to untreated plot) (%)	
	Herbicidal activity	Phytotoxicity
0	61 -	100
1	41 - 60	90 - 99
2	21 - 40	80 - 89
3	11 - 20	60 - 79
4	1 - 10	40 - 59
5	0	0 - 39

The following compounds were used in the Examples for comparison: ●

0075267

Compound No.	Structure	Remarks
(a)		U.S. patent 4,249,934
(b)		Japanese Patent Publn. (un- examined) No. 44587/78
(c)		U.S. patent 4,249,934
(d)		Commercially available herbicide known as "Alachlor"
(e)		Commercially available herbicide known as "Atrazine"
(f)		Commercially available herbicide known as "Ioxynil"


0075267

Compound
No.

Structure

Remarks

(g)



Commercially
available
herbicide
known as "Chlor-
methoxynil"

Test Example 1

Plastic beakers (diameter, 10 cm; height, 10 cm) were filled with upland field soil, and the seeds of barnyardgrass, wild oat, tall morningglory and velvetleaf and the seeds of soyben, cotton and corn were separately sowed in the beakers. A designed amount of the test compound formulated into an emusifiable concentrate according to Formulation Example 2 or 6 and dispersed in water was sprayed over the top by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were grown for 20 days in the greenhouse, and herbicidal activity and phytotoxicity were examined. The results are shown in Table 8.

0075267

Table 8

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity				Phytotoxicity		
		Barn-yard-grass	Wild oat	Tall morning-glory	Velvet-leaf	Soybean	Cotton	Corn
1	5	5	5	5	5	1	1	1
2	2.5	5	5	5	5	1	1	1
3	2.5	5	5	5	5	1	1	1
4	2.5	5	5	5	5	1	1	1
6	5	5	5	5	5	1	1	1
8	2.5	5	5	5	5	1	1	1
9	2.5	5	5	5	5	1	1	1
10	5	5	5	5	5	1	1	1
12	2.5	5	5	5	5	1	1	1
14	2.5	5	5	5	5	1	1	1
16	2.5	5	5	5	5	1	1	1
17	2.5	5	5	5	5	1	1	1
18	2.5	5	5	5	5	1	1	1
(a)	5	1	0	0	3	1	1	0
(b)	2.5	3	1	3	5	2	3	1
(c)	2.5	1	1	0	4	0	0	1

0075267

Test Example 2

In plastic pots (diameter, 10 cm; height, 10 cm) filled with upland field soil, the tubers of purple nutsedge were transplanted and cultivated in a greenhouse for 4 weeks. Separately, the seeds of barnyardgrass, wild oat, wild mustard and velvetleaf were sowed in the similar pots and grown for 2 weeks in the greenhouse. A designed amount of the test compound formulated in an emulsifiable concentrate according to Formulation Example 2 or 6 and dispersed in water with a spreading agent was sprayed to the foliage of the test plants by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were further grown for 3 weeks in the greenhouse, and the herbicidal activity was examined. The results are shown in Table 9.

Among the test plants, purple nutsedge was further grown for 2 weeks in the greenhouse, and the number of revived sprouts in the treated plot was compared with that in the untreated plot. Evaluation was made on the following criteria: A, no revival; B, 1 to 10 % revival; C, 11 - 50 % revival; D, more than 51 % revival. The results are also shown in Table 9.

0075267

Table 9

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity					Prevention of revival
		Barnyard-grass	Wild oat	Wild mustard	Velvet-leaf	Purple nutsedge	
1	5 1.25	4	5	5	5	A	A
2	5 1.25	4	3	5	5	-	-
3	20 5	3	3	5	5	-	-
4	5 1.25	-	4	5	5	-	-
6	5 1.25	5	5	5	5	-	-
8	20 5	2	3	3	5	-	-
9	20 1.25	-	5	5	5	-	-
10	5 1.25	5	5	5	5	-	-
12	20 1.25	2	2	4	5	-	-
14	5 1.25	5	4	4	5	-	-
16	5 1.25	2	2	3	5	-	-
17	5 1.25	3	3	5	5	-	-
18	20 1.25	4	3	5	5	-	-
(a)	5 1.25	0	0	1	4	-	-
(b)	5 1.25	0	0	2	5	-	-
(c)	5 1.25	0	0	0	4	-	-

0075267

Test Example 3

Plastic trays (35 cm x 25 cm x 15 cm) were filled with upland field soil, and the seeds of tall morningglory, velvetleaf, prickly sida, jimsonweed, black nightshade, redroot pigweed, johnsongrass and green foxtail, and the seeds of cotton and soybean were sowed therein. A designed amount of the test compound formulated into a wettable powder according to formulation Example 1 or 5 and dispersed in water was sprayed over the top by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were grown in a greenhouse for 20 days, and phytotoxicity and herbicidal activity were examined. The results are shown in Table 10.

0075267

Table 10

Com-pound No.	Dosage (weight of active in-gredient, g/are)	Herbicidal activity								Phyto-toxicity	
		Tall morning-glory	Velvet-leaf	Prickly sida	Jimson-weed	Black night-shade	Redroot pigweed	Johnson-grass	Green foxtail	Cotton	Soybean
3	5	5	5	5	5	5	5	5	5	1	-
10	2.5	4	5	5	5	5	5	5	5	0	-
	5	5	5	5	5	5	5	5	5	1	1
	2.5	4	5	5	5	5	5	4	5	0	0
(a)	5	0	4	2	0	3	5	5	0	2	0
(b)	5	2	5	4	3	5	5	5	4	3	2
(c)	2.5	0	4	3	1	4	5	5	2	2	0
(d)	5	0	5	2	0	4	5	5	2	0	-
	20	1	0	2	0	4	5	2	5	0	0

0075267

Test Example 4

Plastic trays (35 cm x 25 cm x 15 cm) were filled with upland field soil, and the seeds of corn, wheat, velvetleaf, cocklebur, tall morningglory, common lambsquarters, black nightshade, common chickweed and green foxtail were sowed and grown for 2 to 3 weeks in a greenhouse. Every two trays were placed in a frame (50 cm x 100 cm x 40 cm) and a designed amount of the test compound was sprayed thereover by means of a small hand sprayer. The test plants were further grown for 3 weeks in the greenhouse and herbicidal activity and phytotoxicity were examined. The results are shown in Table 11. In this treatment, the test compound was formulated into an emulsifiable concentrate according to Formulation Example 2 and applied by diluting it in water (25 liters) with the addition of a spreading agent. At the time of application, the plants were generally at the 1 to 4 leaf stage and in 1.5 to 20 cm height.

0075267

Table 11

Com-pound No.	Dosage (weight of active in-gredient, g/are)	Phyto-toxicity		Herbicidal activity						
		Corn	Wheat	Velvet-leaf	Cock-lebur	Tall morning-glory	Common lambs-quarters	Black night-shade	Common chick-weed	Green fox-tail
1	0.63	1	1	5	5	5	5	5	4	3
14	0.32	1	0	5	5	5	5	4	3	2
	0.63	-	1	5	5	5	5	5	4	4
	0.32	1	1	5	5	5	5	5	2	2
(e)	2.5	0	-	5	5	5	-	-	-	1
(f)	1.25	0	-	4	3	3	-	-	-	0
	2.5	-	0	-	-	-	5	5	5	0
	1.25	-	0	-	-	-	5	4	3	0

0075267

Test Example 5

Wagner's pots (1/5000 are) were filled with paddy field soil and the seeds of annual weeds (e.g. barnyard-grass, monochoria, broad-leaved weeds) were sowed to 2 to 3 cm depth, and water was poured therein. The tubers or buds of perennial weeds (e.g. slender spikerush, hardstem bulrush, arrowhead) were transplanted therein and grown for 5 days. At the time when the germination occurred, a desinged amount of the test compound formulated in an emulsifiable concentrate according to Formulation Example 2 or 6 was applied to the pots by perfusion. Thereafter, the test plants were grown for further 4 weeks and herbicidal activity was examined. The results are shown in Table 12.

0075267


Table 12

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity					
		Barnyard-grass	Monochoria	Broad-leaved weed	Slender spikerush	Hardstem bulrush	Arrow-head
9 14	0.5	5	5	5	5	5	5
	0.25	4	5	5	4	4	4
	0.5	5	5	5	5	5	5
	0.25	4	5	5	4	4	4
(a)	0.5	0	3	3	1	1	0
	0.25	0	1	1	0	1	0
(b)	0.5	0	3	3	2	1	0
	0.25	0	2	1	1	0	0
(c)	0.5	0	3	1	1	2	1
	0.25	0	2	1	0	0	0
(g)	0.5	2	2	2	2	1	1
	0.25	1	2	2	1	0	0

0075267

What is claimed is:

1. A condensed phenylurazol of the formula:



wherein X is a chlorine atom or a bromine atom, Y and Z are each an oxygen atom or a sulfur atom and R is a C_1-C_3 alkyl group, an allyl group or a propargyl group and n is an integer of 4 or 5.

2. The condensed phenylurazol according to claim 1, wherein at least one of Y and Z is an oxygen atom.

3. 2-(4-Chloro-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]pyridazine-1,3(2H)-dione.

4. 2-(4-Chloro-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]pyridazine-3-thioxo-1(2H)-one.

5. 2-(4-Chloro-2-fluoro-5-methoxyphenyl)-1H,5H-[1.2.4]triazolo[1.2-a][1.2]diazepine-1,3(2H)-dione.

6. 2-[4-Chloro-2-fluoro-5-(1-methylethoxy)-phenyl]-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]-


0075267

pyridazine-1,3(2H)-dione.


7. 2-(4-Chloro-2-fluoro-5-propoxyphenyl)-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]pyridazine-1,3(2H)-dione.

8. 2-[4-Chloro-2-fluoro-5-(2-propynyloxy)-phenyl]-5,6,7,8-tetrahydro-1H-[1.2.4]triazolo[1.2-a]-pyridazine-1,3(2H)-dione.

9. A process for producing a condensed phenylurazol of the formula:




wherein X is a chlorine atom or a bromine atom, Y and Z are each an oxygen atom or a sulfur atom but at least one of Y and Z is an oxygen atom and R is a C_1-C_3 alkyl group, an allyl group or a propargyl group and n is an integer of 4 or 5, which comprises reacting a phenylurazol of the formula:




wherein X, Y, Z and R are each as defined above in a metallized form with a dihaloalkane of the formula:

0075267




wherein A and A' are each a halogen atom and n is as defined above.

10. A process for producing a condensed phenylurazol of the formula:




wherein X is a chlorine atom or a bromine atom, Y is an oxygen atom or a sulfur atom and R is a C₁-C₃ alkyl group, an allyl group or a propargyl group, which comprises treating a urea of the formula:



wherein R' is a lower alkyl group and X, Y and R are each as defined above with a base.


11. A process for producing a condensed phenylurazol of the formula:



wherein X is a chlorine atom or a bromine atom, Y is an


0075267

oxygen atom or a sulfur atom and R is a C₁-C₃ alkyl group, an allyl group or a propargyl group, which comprises reacting a urea of the formula:




wherein X, Y and R are each as defined above with phosgene.

13. A process for producing a condensed phenylurazol of the formula:




wherein X is a chlorine atom or a bromine atom, Y is an oxygen atom or a sulfur atom and R is a C₁-C₃ alkyl group, an allyl group or a propargyl group, which comprises reacting a urea of the formula:




wherein X, Y and R are each as defined above with carbon disulfide.

14. A process for producing a condensed phenylurazol of the formula:

0075267



wherein X is a chlorine atom or a bromine atom, R is a C_1-C_3 alkyl group, an allyl group or a propargyl group and n is an integer of 4 or 5, which comprises reacting a condensed phenylurazol of the formula:



wherein X, R and n are each as defined above with phosphorus pentasulfide.


15. A herbicidal composition which comprises a herbicidally effective amount of the condensed phenylurazol according to claim 1 and an inert carrier.

16. A method for controlling or exterminating weeds which comprises applying a herbicidally effective amount of the condensed phenylurazol according to claim 1 to the area where the weeds grow or will grow.

17. Use of the condensed phenylurazol according to claim 1 as a herbicide.

0075267

18. A urea of the formula:



wherein X is a chlorine atom or a bromine atom, Y is an oxygen atom or a sulfur atom, R is a C₁-C₃ alkyl group, an allyl group or a propargyl group and R' is a lower alkyl group.



European Patent
Office

EUROPEAN SEARCH REPORT

0075267
Application number

EP 82 10 8507.3

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D,Y	US - A - 4 249 934 (O. WAKABAYASHI et al.) * claims 1 to 4, 13, 14, 18, 19, 23, 24; column 7, line 13 to column 8, line 39 *	1,2, 13,15, 17	C 07 D 487/04 C 07 D 237/04 A 01 N 43/64 /(C 07 D 487/04, 249/00,243/00) (C 07 D 487/04, 249/00, 237/00)
D,A	* columns 2, 3; column 5, line 5 to column 7, line 12 *	9,10, 14	
Y	US - A - 4 214 891 (A.D. WOLF) * claims 1, 2, 4, 5, 7, 8, 10, 11; tables II, III *	1,2, 15,17	TECHNICAL FIELDS SEARCHED (Int.Cl. 3)
A	* claims 13, 14, 16, 17 *	16	C 07 D 237/02 C 07 D 237/04 C 07 D 249/16 C 07 D 487/02 C 07 D 487/04
Y	Patent Abstracts of Japan Vol. 2, No. 85, 12 July 1978 page 1253C78	1,2, 13,15 17	
Y	& JP - A - 53 - 44587		
A	Chemical Abstracts vol. 95, no. 7, 17 August 1981 Columbus, Ohio, USA M.J. KORNET et al. "Synthesis of 1- methyl- and 1-ethoxycarbonyl-2-phenyl- carbamoylpiperidazines as potential anticonvulsant agents" page 712, abstract no. 62107k & Journal of Heterocyclic Chemistry, vol. 18, no. 2, 1981 pages 293 to 295	18	CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document
X The present search report has been drawn up for all claims			
Place of search	Date of completion of the search	Examiner	
Berlin	09-12-1982	HASS	



European Patent
Office

EUROPEAN SEARCH REPORT

0075267
Application Number

EP 82 10 8507.3

- page 2 -

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	Chemical Abstracts vol. 88, no. 7, 13 February 1978 Columbus, Ohio, USA O. WAKABAYASHI et al. " Herbicidal N,N'-alkylene-N-alkoxycarbonyl-N'-(thio) carbamoylhydrazines" page 553, column 2, abstract no. 50904k & JP - A - 77 - 83552 -----	18	
			TECHNICAL FIELDS SEARCHED (Int. Cl.)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.